

## REFERENCES

- Anon (2000). Assessment of impact of offshore wind energy structures on the marine environment. Byrne Ó Cléirigh / Ecoserve report prepared for The marine Institute, Ireland, 44pp.
- Bray, J.R. & Curtis, J.T. (1957) An ordination of the upland forest communities of Southern Wisconsin. *Ecol. Monogr.*, 27, 325-349.
- Buchanan, J.B. (1984) Sediment Analysis In: Holme, N.A & McIntyre, A. Methods for the study of Marine Benthos. Blackwell Scientific Publications, London. pp 41-65.
- Camphuysen, K., Fox, T., Leopold, M., & Petersen, K. (2004) Towards standardised seabirds at sea census techniques in connection with environmental impact assessments for offshore wind farms in the U.K. A Comparison of Ship and Aerial Sampling Methods for Marine Birds, and Their Applicability to Offshore Wind Farm Assessments. Koninklijk Nederlands Instituut voor Onderzoek der Zee Report commissioned for COWRIE.
- Clark, K.R. & Warwick, R.M. (1994) Change in Marine Communities: An Approach to Statistical Analysis and Interpretation. Natural Environment Research Council and Plymouth Marine Laboratory: 144 pp.
- CMACS & MarineSeen (2004) Biology & Video Surveys of North Hoyle Wind Turbines. Internal report for npower renewables.
- CMACS (2002) North Hoyle Offshore Windfarm Environmental Impact Assessment – technical report. Marine Sediments. Centre for Marine and Coastal Studies, University of Liverpool January 2002, report no 2642.
- CMACS (2003). A baseline assessment of electromagnetic fields generated by offshore windfarm cables. COWRIE Report EMF-01-2002, 71pp.
- Connor, D.W., Dalkin, M.J., Hill, T.O., Holt, R.H.F. & Sanderson, W.G. (1997) Marine Nature Conservation Review: marine biotope classification for Britain and Ireland. Volume 2. Sublittoral biotopes. Version 97.06. JNCC Report, No. 230.
- Cooper, W. (2005) North Hoyle Offshore Wind Farm: Review of Scour. ABP Marine Environmental Research Ltd.
- Cranfield University, CMACS Ltd & CEFAS (2006) COWRIE 2: Electromagnetic Fields (EMF) Phase 2. Stage 1 Project Plan. COWRIE-EMF-1-06
- Danish Institute for Fisheries Research (2002). Effects of marine windfarms on the distribution of fish, shellfish and marine mammals in the Horns Rev area. DFU-rapport 117-02, 42pp.
- Ellis, J.R. and Parker-Humphreys, M. (2004). Fish populations in the eastern Irish Sea. Report to National Wind Power, C2150/01, 29pp.
- Ellis, J.R. and Parker-Humphreys, M. (2005). Fish populations in the eastern Irish Sea. Report to Npower Renewables, C2150/02, 28pp.
- Ellis, J.R., Armstrong, M.J, Rogers, S.I., and Service, M. (2002). The distribution, structure and diversity of fish assemblages in the Irish Sea. In Marine biodiversity in Ireland and adjacent waters (J.D.Nunn, ed.). Ulster Museum, Belfast, 93–114.

Ellis, J.R., Rogers, S.I. and Freeman, S.M. (2000). Demersal assemblages in the Irish Sea, St George's Channel and Bristol Channel. *Estuarine and Coastal Shelf Science*, 51(3):299–315.

Folk, R.L. 1954. The distinction between grain size and mineral composition in sedimentary-rock nomenclature. *Journal of Geology*, 62, 344-359.

Fugro-UDI Ltd, 2001. Survey Services Report, North Hoyle Wind Farm Geophysical Survey.

Gill, A.B. and Taylor, H. (2001). The potential effects of electromagnetic fields generated by cabling between offshore wind turbines upon elasmobranch fishes. CCW Science Report No, 488, 73pp.

Gislason, K (2003) Evaluation of Scour. LIC Memo 0148-53.

Hammond, P.S, Northridge, S.P., Thompson, D., Gordon J.C.D., Hall, A.J., Aarts G. and Matthiopoulos, J. (2005) Background information on marine mammals for Strategic Environmental Assessment 6. Sea Mammal Research Unit, St Andrews.

Hayward, P.J & Ryland, J.S. (1994) *The Marine Fauna of the British Isles and North West Europe*. Oxford University Press. Pp951.

Howson, C.M. & Picton, B.E. (eds) (1997) *The species directory of the marine fauna and flora of the British Isles and surrounding seas*. Ulster Museum and Marine Conservation Society, Belfast and Ross-on-Wye. Ulster Museum Publication No. 276.

Hyperbaric Consult A/S (2003) Scour Survey Note. Report HBC-750-NH-R002.

Innogy plc (2002). North Hoyle Offshore Wind Farm: Environmental Statement. NWP Offshore Ltd.

Lockwood, S.J. (2005) Gwynt y Môr Offshore Wind Farm Technical Report: Commercial & Recreational Fisheries of Liverpool Bay. A report prepared on behalf of npower renewables Ltd.

May, J. (2005) Overview of Sea Bed Bathymetric Levels between 2001 and 2004 (Pre and Post Construction). PMSS Ltd.

Moore, J. (2002) An atlas of marine biodiversity action plan species and habitats and species of conservation concern in Wales. 2nd Edition. CCW contract science report no 509.

Npower renewables (2005a) North Hoyle Offshore Wind Farm FEPA Monitoring Report. Npower renewables.

Npower renewables (2006a) North Hoyle Offshore Wind Farm FEPA Monitoring Report (2004-5). Npower renewables.

NWPO (2003) Project Method Statement. Rev 2.

Osiris Projects Ltd (2006) North Hoyle Offshore Wind Farm. Post construction Geophysical Survey Report. Data Collection Period - 5th and 16th April 2006

Osiris Projects Ltd (2005) North Hoyle Offshore Wind Farm. Post construction Geophysical Survey Report. Data Collection Period - 12th August –12nd October 2004. C4014.

Osiris Projects Ltd (2005) North Hoyle Offshore Wind Farm. Scour Monitoring Surveys Report. Data Collection Period - 26th April –02nd May 2005. C5011.

Parker-Humphreys, M. (2004). Distribution and relative abundance of demersal fishes from beam trawl surveys in the Irish Sea (ICES Division VIIa) 1993–2001. Science Series Technical Report, CEFAS, Lowestoft, 120, 68pp.

Rees, E.I.S. (2001) Habitat specialization by *Thia scutellata* (Decapoda: Brachyura) off Wales. Journal of the Association of the marine Biological Association of the United Kingdom. 81. pp 697-698.

Reid, J., Evans, P.G.H. and Northridge, S.P. (2003) Cetacean Distribution Atlas. Joint Nature Conservation Committee, Peterborough. 68pp.

Rigden, D. (2004). UK wind to generate over 7GW under new initiatives. Marine Scientist, 6, 34–36.

Stone, C.J., Webb, A., Barton, C., Ratcliffe, N., Reed, T.C., Tasker, M.L. Camphuysen, C.J. & Pienkowski, M.W. (1995) An atlas of seabird distribution in north-west European waters. Joint Nature Conservation Committee and Nederlands Instituut voor Onderzoek der Zee, Peterborough.

Thomsen, F., Ludemann, K., Kafemann, R. & Piper, R (2006) Effects of Offshore Wind Farm Noise on Marine Mammals and Fish. Biola, Hamburg, Germany on behalf of COWRIE Ltd.

Tougaard, J., Carstensen, J., Henriksen, O.D., Skov, H., and Teilmann, J. (2003) Short-term effects of the construction of wind turbines on harbour porpoises at Horns reef. Technical report to TechWise A/S. HME/362-02662. Hedeselskabet, Roskilde, Denmark.

Wheeler, A. (1969). The fishes of the British Isles and North West Europe. Michigan State University Press, 613pp.

Wheeler, A. (1978). Key to the Fishes of Northern Europe. Frederick Warne, London. 380pp.

Whitehead, P.J.P., Bauchot, M.L., Hureau, J.-C., Nielsen, J. and Tortonese, E. (Eds.) (1984). Fishes of the North-eastern Atlantic and the Mediterranean, Vol. 1–3. UNESCO, Paris, 1473pp.

WWT (2005) Aerial Surveys of Waterbirds in Strategic Windfarm Areas: 2004/05 Final Report. WWT Wetlands Advisory Service Report to Department of Trade and Industry, October 2005